

Species	States Where Sensitive (a)				Forests Where Species is Known (K) or Suspected (S) to Occur (b)														Comments
	MT	ID	ND	SD	B/D	BRT	CL W	CUS	DPG	FLA T	GAL	HEL	IPN F	KOOT	L&C	LOLO	NEZ		
FISH																			
Pacific Lampry ( <i>Lampetra tridentata</i> )		S1					S	K										K	
Northern Redbelly Dace ( <i>Phoxinus eos</i> )			S2	S2					S	K									
*Sturgeon Chub ( <i>Macrhybopsis gelida</i> )	S2		S2	S2					K	E/S								E/S - extirpated or suspected	
*Yellowstone Cutthroat ( <i>Oncorhynchus clarki bouvieri</i> )	S2	S2							K			K							
*Fluvial Arctic Grayling ( <i>Thymallus arcticus montanus</i> )	S1					K						S				S			
*Westslope Cutthroat ( <i>Oncorhynchus clarki lewisi</i> )	S2	S3				K	K	K			K	K	K	K	K	K	K	K	
*Interior Redband ( <i>Oncorhynchus mykiss gairdneri</i> )	S1	S4						S					K	K				K	
*Snake River SPR/SUM Chinook ( <i>Oncorhynchus tshawtscha</i> )		S1						K										K	
*Burbot ( <i>Lota lota</i> )		S1											K					Northern Idaho only	

\* Indicates this species was on the 1999 USFS R1 Sensitive Species List

(a) Species are listed as Sensitive by State. The State where a species is listed as Sensitive is indicated by an "X" in the State/species column. A species identified as Sensitive within a State, will be considered as Sensitive on all Units within the State where it occurs, unless described otherwise.

(b) National Forest (Grasslands) where a species is known or suspected to occur, within States where a species is listed as Sensitive, are identified by shading and either a known "K" or suspected "S" in the Forest/species column